

Generalized Barrett-Crane Vertices and Invariants of Embedded Graphs

David N. Yetter
 Department of Mathematics
 Kansas State University
 Manhattan, KS 66506

1

1 Introduction

In [1] Barrett and Crane introduce a modification of the generalized Crane-Yetter state-sum (cf. [2]) based on the category of representations of $Spin(4) \cong SU(2) \times SU(2)$, which provides a four-dimensional analogue of Regge and Ponzano's [9] spin-network formulation of three-dimensional gravity.

The key to the modification of the Crane-Yetter state-sum is the use of a different intertwiner between the “inbound” and “outbound” tensor products of objects assigned to faces of the tetrahedron, thereby imposing a “quantum analogue” of the condition that the sum of the simple bivectors represented by two faces with a common edge is itself simple. The purposes of this paper are

1. to give an explicit formula for the Barrett-Crane intertwiners and their q -analogues by using the Kauffman-Lins [6] “Temperley-Lieb” recoupling theory for (quantized) $SU(2)$ in each factor,
2. to show that in the case where the $SU(2)$ factors of $Spin(4)$ are deformed with reciprocal deformation parameters the operators have the symmetry properties of a topological 4-vertex when represented in the usual diagrammatic calculus for braided tensor categories (cf. [5], [4] [11]),
3. to generalize the family of operators to families which exhibit the symmetry properties of topological n -vertices when represented in the diagrammatic calculus, and
4. to consider the natural family of invariants of topological graphs embedded in $\mathbb{R}^3$ or S^3 which arise by applying the functorial construction first proposed in [4] and [12].

Although the present paper will be concerned with the construction of “algebraic n -vertices” and their applications to topological graph theory, the operators constructed herein may well have applications to the physically-motivated Barrett-Crane state-sum, in particular allowing computations of state spaces and transition amplitudes from cell-decompositions coarser than triangulations.

¹Supported by NSF Grant # DMS-9504423

2 $Spin(4)$ Recoupling Theory via Temperley-Lieb Recoupling


The factorization $Spin(4) \cong SU(2) \times SU(2)$ (or rather $\mathfrak{so}_4 \cong \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$) has an easy, immediate consequence for the structure of the associated QUEA: the standard 1-parameter deformation $U_q(\mathfrak{sl}_2)$ gives rise to a 2-parameter deformation $U_{q,r}(\mathfrak{so}_4) \cong U_q(\mathfrak{sl}_2) \otimes U_r(\mathfrak{sl}_2)$, where q is the deformation parameter for the first $\mathfrak{sl}_2$, and r for the second.

We will be interested in 1-parameter deformation which arises by specializing to $r = q^{-1}$, which we will denote $U_q^{\text{bal}}(\mathfrak{so}_4)$. The irreducible representations of $U_q^{\text{bal}}(\mathfrak{so}_4)$ are each a tensor product of an irreducible representation of $U_q(\mathfrak{sl}_2)$ with an irreducible representation of $U_{q^{-1}}(\mathfrak{sl}_2)$, and thus are indexed by pairs of non-negative half-integers (spins), or more conveniently, since we will be using Kauffman-Lins style recombination by integers (twice spin = dimension -1 = number of strands q -symmetrized).

Throughout the following we adopt the notation of Kauffman-Lins [6] for all aspect of $U_q(\mathfrak{sl}_2)$ recoupling theory. It should be observed that it will be unnecessary to have different notations for $U_q(\mathfrak{sl}_2)$ and $U_{q^{-1}}(\mathfrak{sl}_2)$ recoupling constants, since the q -symmetrizer, three-vertex, quantum dimensions, $\theta(a, b, c)$, and q -6j symbols are all unchanged when q is replaced with q^{-1} .

We can then let a recombination network labelled with pairs of twice-spins represent the tensor product of the corresponding network labelled with first twice-spins with that labelled with second twice-spins, preceeded and followed by the necessary coherence maps from the underlying category of vector-spaces.

For example



represents the map

$$(V_i \otimes W_j) \otimes (V_k \otimes W_l) \xrightarrow{c} (V_i \otimes V_k) \otimes (W_j \otimes W_l) \xrightarrow{v_{ikm} \otimes w_{jln}} V_m \otimes W_n$$

where c is the middle-four-interchange map composed of associators and symmetrizers in $\mathbb{C} - v.s.$, and v_{ikm} (resp. w_{jln}) is the map named by the Kauffman-Lins three-vertex with the chosen labels.


By the same reasoning, it follows that the quantum dimension of (i, j) is $\Delta_i \Delta_j$, and similarly that the evaluation of a closed diagram labelled with pairs

of twice-spins is the product of the evaluations of the corresponding diagrams obtained by selecting all of the first (resp. second) twice-spins as labels. (In the case where there are braidings, one must use the correct deformation parameter in each case).

A crucial ingredient in the Barrett-Crane state sum and in what follows is the notion of a “balanced” irreducible, that is one of the form (j, j) for some j . Several facts stand out about balanced irreducibles in recombination diagrams:

First, it should be observed that a curl on a strand with a balanced label is simply the identity operator: the constants contributed by each tensor and cancel (because we are working with $U_q^{\text{bal}}(\mathfrak{so}_4)$).

Second, and for the same reason, we have




Thus we see that for balanced irreducibles, the curl is trivial, while the (family of) 3-vertex admits as symmetries all permutations, not just cyclic permutations as in Kauffman-Lins (or for non-balanced labels in the present setting).

We are now in a position to introduce the formula in Kauffman-Lins notation for the Barrett-Crane 4-vertex.

3 Barrett-Crane Vertices, Properties and Generalizations

Definition 3.1 *The Barrett-Crane 4-vertex $v_{k,l}^{i,j}$ is the map from $V_{(i,i)} \otimes V_{(j,j)}$ to $V_{(k,k)} \otimes V_{(l,l)}$ given by the sum of recombination networks*



From the point of view of this paper the crucial properties of these map are their symmetry properties. It is immediate from the observations above and the cyclic symmetry properties of 3-vertices (which follow from the corresponding properties for Kauffman-Lins 3-vertices) that

$$\sigma_{(j,j),(i,i)}(v_{k,l}^{i,j}) = v_{k,l}^{j,i}$$

$$v_{k,l}^{i,j}(\sigma_{(k,k),(l,l)}) = v_{l,k}^{i,j}$$

$$\epsilon_{(i,i) \otimes (j,j)} \otimes V_{(l,l)} \otimes V_{(k,k)} (V_{(j,j)} \otimes V_{i,i} \otimes v_{k,l}^{i,j} \otimes V_{(l,l)} \otimes V_{(k,k)} (V_{(j,j)} \otimes V_{(i,i)} \otimes \eta_{(k,k) \otimes (l,l)})) = v_{j,i}^{l,k}$$

and

$$V_{(l,l)} \otimes V_{(k,k)} \otimes \epsilon_{(i,i) \otimes (j,j)} (V_{(l,l)} \otimes V_{(k,k)} \otimes v_{k,l}^{i,j} \otimes V_{(j,j)} \otimes V_{i,i} (\eta_{(k,k) \otimes (l,l)} \otimes V_{(j,j)} \otimes V_{(i,i)})) = v_{j,i}^{l,k}$$

where $\eta(-)$ and $\epsilon(-)$ are the duality transformations in $Rep(U_q^{\text{bal}}(\mathfrak{so}_4))$ normalized so that they are given by the “cup” and “cap” networks, and intervening generalized associators have been omitted by Mac Lane’s coherence theorem [8].

What is remarkable is that the Barrett-Crane 4-vertex can also be expressed in terms of the same formula with the network rotated by $\pi/2$, that is:

Proposition 3.2

$$v_{k,l}^{i,j} = \epsilon_{(k,k)} \otimes V_{(i,i)} \otimes V_{(j,j)} (V_{(k,k)} \otimes v_{l,j}^{k,i} \otimes V_{(j,j)} (V_{(k,k)} \otimes V_{(l,l)} \otimes \eta_{(j,j)}))$$

where ϵ and η are as above, and the intervening associators have been omitted.


proof: The calculation with linear combinations of recombination networks is given in Figure 1, where $\delta_{r,s}$ is the Kronecker delta. The first equation follows from the fact that the evaluation of a doubled planar network (including all those in the Kauffman-Lins construction of a q-6j symbol) is simply the product of the two corresponding $U_q(\mathfrak{sl}_2)$ networks, the second by the symmetry properties of the q-6j symbol, the third is elementary, the fourth by the orthogonality properties of the q-6j symbols. $\square$

We will also refer to maps obtained from Barrett-Crane 4-vertices by tensoring $V_{k,l}^{i,j}$ on the right (resp. right, left, left) with (j,j) (resp. (l,l) , (i,i) , (k,k)) and pre- (resp. post-, pre-, post-)composing with $V_{(i,i)} \otimes \eta_{(j,j)}$ (resp. $V_{(k,k)} \otimes \epsilon_{(l,l)}$, $\eta_{(i,i)} \otimes V_{(j,j)}$, $\epsilon_{(k,k)} \otimes V_{(l,l)}$) as Barrett-Crane 4-vertices, as well as those obtained by tensoring $V_{k,l}^{i,j}$ on the right (resp. right, left, left) with (j,i) (resp. (l,k) , (j,i) , (l,k)) and pre- (resp. post-, pre- post-)composing with $\eta_{(i,i) \otimes (j,j)}$ (resp. $\epsilon_{(l,l) \otimes (k,k)}$, $\eta_{(j,j) \otimes (i,i)}$, $\epsilon_{(k,k) \otimes (l,l)}$). Less formally, but more intelligibly, the maps just described are those obtained by using duality to “turn” some of the inputs or outputs of the vertex “down” or “up” – the geometry of the spin-networks coinciding nicely with raising and lowering of indices.

It thus follows from Proposition 3.2 and the symmetry properties already observed any map in $Rep(U_q^{\text{bal}}(\mathfrak{so}_4))$ between tensor products of balanced objects which admits an expression in terms of a connected recombination network containing a single Barrett-Crane 4-vertex is itself a Barrett-Crane 4-vertex in this more general sense.


Before generalizing to n -vertices, it is convenient to change notation slightly. Let an unmarked node where three edges labelled (i,i) , (j,j) and (k,k) meet denote $\frac{1}{\theta(i,j,k)}$ times the 3-vertex denoted with the heavy dot:

In this new notation the formula for a Barrett-Crane 4-vertex becomes



With this change of notation, it then becomes clear how to define n -vertices for any $n \geq 3$.

For $n = 3$ the 3-vertex is the “new 3-vertex”



for $n = 4$ it is the Barrett-Crane 4-vertex already defined, while for $n > 4$, an n -vertex with given tensor product of k balanced irreducibles as source, and given tensor product of $n - k$ balanced irreducibles as target can be described by a sum of recombination networks with underlying graph a fixed tree with n leaves (or better still n “external edges” with a free end incident with a vertex) divided into a set of k at the top, and a set of $n - k$ at the bottom and all internal vertices trivalent, with the external edges labelled with the tensorands of the source and target with the, summed over all admissible *balanced* labellings of the internal edges with coefficients equal to the product $\prod_{k=1}^{n-2} \Delta_{j_k}$, where the $n - 2$ internal edges are labelled $(j_1, j_1), \dots, (j_{n-2}, j_{n-2})$, and the $n - 1$ internal vertices are all evaluated a “new 3-vertices”.

An example is shown in Figure 2.

It is then easy to apply Proposition 3.2 to show

Theorem 3.3 *The value of an n -vertex depends only on the source and the target.*

proof: Given any two trivalent trees with the same source and target leaves, we can obtain one from the other by iteratively “fusing” edges as in Proposition 3.2. $\square$

Once this is known, it is trivial to apply the observation that three-vertices with balanced labels absorb braidings and are rotated by duality maps to justify the name of n -vertex for these operators. In particular, recalling that a prolongation of a map in a tensor category is an arbitrary tensor product of the map with identity maps, we have:

Theorem 3.4 *Any composition of a prolongation of an n -vertex with prolongations of braidings, (associators, unit transformations,) and duality transformations which admits a description as a connected and simply-connected recombination diagram when the n -vertex is represented as a single n -valent vertex is itself an n -vertex.*

sketch of proof: Use the coherence theorem for tortile categories [10] to rewrite all braidings and unit transformations as compositions of prolongations of braidings, respectively unit transformations, indexed by single (balanced) irreducibles, not tensor products. For each braiding or duality transformation composed directly with the n -vertex, use the previous proposition to express the n -vertex in terms of a linear combination of trees so that all objects indexing the braiding or unit transformation correspond to arcs incident with a single trivalent vertex of the tree. The braiding or duality transformation may then be removed by applying the symmetry properties for 3-vertices noted above. $\square$

What is not immediately clear is that there are also good canonical choices for 0-, 1-, and 2-vertices. Observe that it is clear that a 0 vertex must be an endomorphism of $(0,0)$, that is a scalar. In considering possible relations with previously defined graph invariants, it will be convenient to choose 1. For 1- and 2-vertices, however, an additional structure present on the n -vertices already constructed suggests a canonical choice in each case.

Proposition 3.5 *If $v_{x_1, \dots, x_k, r}^{y_1, \dots, y_{n-k-1}}$ denotes an n -vertex for $n \geq 3$ with source $V_{(y_1, y_1)} \otimes \dots \otimes V_{(y_{n-k-1}, y_{n-k-1})}$ and target $V_{(x_1, x_1)} \otimes \dots \otimes V_{(x_k, x_k)} \otimes V_{(r, r)}$, then*

$$\sum_r \Delta_r v_{x_1, \dots, x_k, r}^{y_1, \dots, y_{n-k-1}} (V_{(x_1, x_1)} \otimes \dots \otimes V_{(x_k, x_k)} \otimes v_{i, j}^r)$$

is an $n+1$ -vertex.

proof: Immediate from the construction of n -vertices. $\square$

If we wish the result to hold without the restriction $n \geq 3$, we must define 2- and 1-vertices by

$$v_j^i = \begin{cases} \frac{1}{\Delta_j} Id_{V_{(j, j)}} & \text{if } i = j \\ 0 : V_{(i, i)} \rightarrow V_{(j, j)} & \text{otherwise} \end{cases}$$

and

$$v_j = \begin{cases} Id_{V_{(0,0)}} & \text{if } i = j \\ 0 : V_{(0,0)} \rightarrow V_{(j,j)} & \text{otherwise} \end{cases}$$

respectively.

The check that these lead to the desired properties is easy and left to the reader.

Thus we see that the full sub-category of tensor products of balanced irreducibles in $Rep(U_q^{bal}(\mathfrak{so}(4)))$ is a “graphical category” in the terminology of [12].

4 Invariants of Embedded Graphs

Some preliminaries are in order:

Definition 4.1 A graph Γ is a pair of sets $V(\Gamma), E(\Gamma)$ whose elements are called vertices and edges respectively, together with a function

$$i_\Gamma : E(\Gamma) \rightarrow V(\Gamma)_1 \coprod V(\Gamma)_2.$$

where S_i denotes the set of i -element subsets of S . Usually elements of the target is thought of as unordered pairs of elements of $V(\Gamma)$. By abuse of terminology and notation, we identify Γ with its geometric realization, the topological space

$$E(\Gamma) \times [0, 1] / \equiv$$

where $\equiv$ is an equivalence relation induced by choosing for each $e \in E(\Gamma)$ a surjection $k_e : \{0, 1\} \rightarrow i_\Gamma(e)$, and defining $(e, x) \equiv (\eta, \xi)$ if and only if they are equal or both x and ξ are 0 or 1 and $k_e(x) = k_\eta(\xi)$.

Observe first that for any two choices of $\equiv$ there is a homeomorphism between the resulting spaces which maps each $\{e\} \times [0, 1]$ to itself either by the identity map or by $t \mapsto 1 - t$, and second that a graph has a natural PL-structure.

Definition 4.2 An embedded graph is a PL-embedding of a graph into $\mathbb{R}^3$. Two embedded graphs are equivalent if they have isomorphic underlying graphs and there exist a PL ambient isotopy of one to the other which preserves the set of vertices.

It is fairly easy to see how to use the family of n -vertices constructed above to construct invariants of embedded graphs: choose a balanced irreducible to label all the edges of the graph and a generic projection of the graph, the invariant is obtained by evaluating the recombination network obtained by interpreting the graphical vertices of valance n as n -vertices in the algebraic sense, crossings as braiding, and maxima and minima as duality transformations. Isotopy invariance is readily verified using the symmetry properties already established for n -vertices and the coherence theorem of Shum [10] for tortile categories.

More generally, we may consider invariants of edge-colored graphs obtained by choosing a balanced irreducible for each color and labelling all edges of that color with the chosen irreducible. In the following, however, we will confine ourselves to invariants obtained by labelling all edges with the same balanced irreducible.

Definition 4.3 Let $\mathcal{G}_j[\Gamma](q)$ denote the value of the invariant obtained by labelling all edges of an embedded graph Γ with the irreducible (j, j) .

We can then easily show

Proposition 4.4 *For j odd, $\mathcal{G}_j[\Gamma](q) \neq 0$ implies Γ is an Eulerian graph, that is each component of Γ is either a single vertex, or admits an Eulerian cycle (equivalently all valences are even).*

proof: Any odd valence vertex will be mapped to an algebraic n -vertex with all incoming labels (j, j) . The underlying Kauffman-Lins 3-vertices in any expression for the n -vertex have value 0. $\square$

Proposition 4.5 *If Γ' is obtained from Γ by reversing all crossings in a regular projection of Γ , then*

$$\mathcal{G}_j[\Gamma'](q) = \mathcal{G}_j[\Gamma](q)$$

proof: Now, $\mathcal{G}_j[\Gamma](q)$ can be computed by fusing arcs exiting each crossing (as in Kauffman and Lins [6]) and summing over all (not necessarily balanced) labellings of the new arcs. Using Kauffman and Lins [6], we may rewrite this sum as a linear combination of evaluations of planar networks at the cost of multiplying each summand by $A^{-k(k+2)+l(l+2)}$ (resp. $A^{k(k+2)-l(l+2)}$) whenever an arc labelled (k, l) occurs at the exit of a positive (resp. negative) crossing. Now observe that since the value of a planar network is unchanged by replacing q by q^{-1} , that given any choice of labels for the new arcs, the value of the summand in the expression for $\mathcal{G}_j[\Gamma]$ corresponding to this choice is equal to the value of the summand in the expression for $\mathcal{G}_j[\Gamma']$ in which the first and second coordinates of each label have been interchanged. Since interchanging coordinates of labels is an involution on the set of labellings of the new arcs, the desired result follows. $\square$

Now, letting $J[L](q)$ denote the Jones polynomial of a link L we have

Proposition 4.6 *If Γ is a disjoint union of cycles, let L be the underlying link obtained by forgetting the vertices, equipped with any orientation, and v_Γ be the number of vertices of Γ , then*

$$\mathcal{G}_1[\Gamma](q) = \frac{J[L](q)J[L](q^{-1})}{\Delta_1^{v_\Gamma}}$$

proof: The value of the $(1,1)$ -labelled network is $J_f[L](q)J_f[L](q^{-1})$ where J_f denotes the framed Jones polynomial (Kauffman bracket, but written in q rather than A). But the normalization constants cancel, so this is the desired numerator. The definition of the 2-vertex contributes a factor of $\frac{1}{\Delta_1}$. $\square$

A standard notion in graph theory is that of a *cutpoint*, that is a vertex whose removal increases the number of connected components. This notion is usually defined in a purely combinatorial way so that the presence of loops at a vertex does not imply that the vertex is a cutpoint. We will need a stronger notion of cutpoint which takes into account the topology and embedding of the graph. First let us recall a standard notion from link theory (in a form applicable to embedded graphs):

Definition 4.7 An embedded graph Γ is a separated union if it is ambient isotopic to an embedded graph $\eta : \Gamma_1 \coprod \Gamma_2 \rightarrow \mathbb{R}^3$ where the images of Γ_1 and Γ_2 lie in disjoint open balls.

Similarly we make

Definition 4.8 An embedded graph Γ is an almost-separated union of two subgraphs Γ_1 and Γ_2 if the intersection of Γ_1 and Γ_2 is a single vertex v , and Γ is ambient isotopic to a graph in which the images of $\Gamma_1 \setminus \{v\}$ and $\Gamma_2 \setminus \{v\}$ lie in disjoint open balls.

Note that here we must regard graphs as their geometric realization, since by $\Gamma_i \setminus \{v\}$ we mean the space obtained by deleting the point v , not the graph obtained by deleting v and all incident edges.

Definition 4.9 A topological cutpoint of an embedded graph Γ is a vertex v such that Γ is an almost-separated union of two subgraphs Γ_1 and Γ_2 with common vertex v . By a splitting of a graph at a topological cut-point, we mean the separated union Γ' of Γ_1 and Γ_2 .

Theorem 4.10 If v is a topological cutpoint of Γ and $\Gamma' \cong \Gamma_1 \coprod \Gamma_2$ is a splitting of Γ at v , then

$$\mathcal{G}[\Gamma] = \mathcal{G}[\Gamma'] = \mathcal{G}[\Gamma_1]\mathcal{G}[\Gamma_2]$$

proof: Observe that if v is a topological cutpoint, we may construct the tree over which the algebraic vertex assigned to v is defined in such a way that there is a distinguished edge whose removal separates Γ_1 and Γ_2 . Further observe that if we use this tree to describe the algebraic vertex at v , and render the evaluation of $\mathcal{G}[\Gamma]$ as a sum of recombination diagrams, all of the diagrams have turn-arounds incident with distinguished edge. By one of the properties of q -symmetrizers noted in [6], it follows that only those summands in which the label on the distinguished edge is $(0, 0)$ can be non-zero.

It may readily be verified that the value of the summand in a algebraic vertex with label $(0, 0)$ on one edge in the tree describing the algebraic vertex is a tensor product of two algebraic vertices whose trees are those obtained by removing the distinguished edge. The result then follows from the lemma below. $\square$

Lemma 4.11 $\mathcal{G}_j$ is multiplicative under separated union.

proof: Immediate from the functorial construction. $\square$

5 Conclusions

The preceding lemma and Proposition 4.4 are strongly suggestive that (for odd j) $\mathcal{G}_j$ may well be related to the Martin polynomial (cf. [7], [3]), at least at $q = \pm 1$ where the braiding is trivial. It would seem to be a fruitful undertaking to examine the invariants described herein in terms of the Hopf algebra structure on the space of linear combinations of embedded graphs induced by the formulas given in [3] for (non-embedded) graphs.

References

- [1] Barrett, J.W. and Crane, L., “Relativistic Spin Networks and Quantum Gravity,” gr-qc 9709028 (to appear CQG).
- [2] Crane, L., Kauffman, L.H., and Yetter, D.N., “State-Sum Invariants of 4-Manifolds,” *JKTR* **4** (2) (1997) 177-234.
- [3] Ellis-Monaghan, J.A., “New Results for the Martin Polynomial,” preprint (1997).
- [4] Freyd, P.J. and Yetter, D.N., “Braided Compact Closed Categories with Applications to Low-Dimensional Topology,” *Adv. in Math.* **77** (2) (1989) 156-182.
- [5] Joyal, A. and Street, R., “The Geometry of Tensor Calculus, I,” *Adv. in Math.* **88** (1) (1991) 55-112.
- [6] Kauffman, L.H. and Lins, S.L., *Temperley-Lieb Recoupling Theory and Invariants of 3-Manifolds*, Princeton Univ. Press (1994).
- [7] Martin, P. *Enumerations Euleriennes dans les Multigraphs et Invariants de Tutte-Grothendieck*, Thesis, Grenoble, 1977.
- [8] Mac Lane, S. *Categories for the Working Mathematician*, Springer (1971).
- [9] G. Ponzano and T. Regge, “Semiclassical Limit of Racah Coefficients” in *Spectroscopic and Group Theoretic Methods in Physics* (R. Bloch, ed), North Holland, Amsterdam, 1968.
- [10] Shum, M.-C. “Tortile Tensor Categories,” *J. Pure and App. Alg.* **93** (1) (1994), 57-110.
- [11] Yetter, D.N., “Quantum Groups and Representations of Monoidal Categories,” *Math. Proc. Camb. Phil. Soc.* **108** (1990) 261-290.
- [12] Yetter, D.N., “Category Theoretic Representations of Embedded Graphs in $\mathbf{S}^3$,” *Adv. in Math* **77** (2) (1989) 137-155.

$$\begin{aligned}
& \sum_n \frac{\Delta_n}{\theta(m.l.n)\theta(j.k.n)} \quad \begin{array}{c} j,j \quad k,k \\ \diagdown \quad \diagup \\ \quad n,n \\ \diagup \quad \diagdown \\ l,l \quad m,m \end{array} \\
&= \sum_{n,r,s} \frac{\Delta_n}{\theta(m.l.n)\theta(j.k.n)} \left\{ \begin{matrix} m & l & r \\ j & k & n \end{matrix} \right\} \left\{ \begin{matrix} m & l & s \\ j & k & n \end{matrix} \right\} \quad \begin{array}{c} j,j \quad k,k \\ \diagdown \quad \diagup \\ \quad r,s \\ \diagup \quad \diagdown \\ l,l \quad m,m \end{array} \\
&= \sum_{n,r,s} \frac{\Delta_n}{\theta(m.l.n)\theta(j.k.n)} \frac{\Delta_r}{\Delta_n} \frac{\theta(m.l.n)\theta(j.k.n)}{\theta(j.l.r)\theta(m.k.r)} \\
& \quad \left\{ \begin{matrix} l & j & n \\ k & m & r \end{matrix} \right\} \left\{ \begin{matrix} m & l & s \\ j & k & n \end{matrix} \right\} \quad \begin{array}{c} j,j \quad k,k \\ \diagdown \quad \diagup \\ \quad r,s \\ \diagup \quad \diagdown \\ l,l \quad m,m \end{array} \\
&= \sum_{n,r,s} \frac{\Delta_r}{\theta(j.l.r)\theta(m.k.r)} \left\{ \begin{matrix} l & j & n \\ k & m & r \end{matrix} \right\} \left\{ \begin{matrix} m & l & s \\ j & k & n \end{matrix} \right\} \quad \begin{array}{c} j,j \quad k,k \\ \diagdown \quad \diagup \\ \quad r,s \\ \diagup \quad \diagdown \\ l,l \quad m,m \end{array} \\
&= \sum_{r,s} \frac{\Delta_r}{\theta(j.l.r)\theta(m.k.r)} \delta_{r,s} \quad \begin{array}{c} j,j \quad k,k \\ \diagdown \quad \diagup \\ \quad r,s \\ \diagup \quad \diagdown \\ l,l \quad m,m \end{array}
\end{aligned}$$

Figure 1: Proof of Proposition 3.2




Figure 2: A 6-vertex